Version with markings to show changes made

- 132. (Amended) A method for inducing acetylcholine receptor synthesis in a cell, comprising contacting said cell with a polypeptide which comprises an epidermal growth factor-like domain, said domain comprising an amino acid sequence set forth in [selected from the group] consisting of SEQ ID NO: 152 [151, 152, 188, 189, and 190], or said domain comprising an amino acid sequence encoded by a nucleic acid sequence selected from the group consisting of [SEQ ID NO: 154,] SEQ ID NO: 155, SEQ ID NO: 156, SEQ ID NO: 157, SEQ ID NO: 158, and SEQ ID NO: 159, wherein said polypeptide is administered in an amount sufficient to stimulate synthesis of acetylcholine receptors in said cell.
- 138. (Twice Amended) The method of claim 132, wherein said epidermal growth factor-like domain comprises an amino acid sequence encoded by a nucleic acid sequence selected from the group consisting of [SEQ ID NO: 154,] SEQ ID NO: 155, SEQ ID NO: 156, SEQ ID NO: 157, SEQ ID NO: 158, and SEQ ID NO; 159.
- 141. (Amended) The method of claim 132, wherein said epidermal growth factor-like domain comprises an amino acid sequence set forth in [SEQ ID NO: 151,] SEQ ID NO: 152 [, or SEQ ID NO: 190].
 - 142. The method of claim 132, wherein said polypeptide binds the p185^{cr6B2} receptor.
 - 143. The method of claim 132, wherein said polypeptide is a recombinant polypeptide.

Pending Claims

- 132. A method for inducing acetylcholine receptor synthesis in a cell, comprising contacting said cell with a polypeptide which comprises an epidermal growth factor-like domain, said domain comprising an amino acid sequence set forth in SEQ ID NO: 152, or said domain comprising an amino acid sequence encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 155, SEQ ID NO: 156, SEQ ID NO: 157, SEQ ID NO: 158, and SEQ ID NO: 159, wherein said polypeptide is administered in an amount sufficient to stimulate synthesis of acetylcholine receptors in said cell.
- 138. The method of claim 132, wherein said epidermal growth factor-like domain comprises an amino acid sequence encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO: 155, SEQ ID NO: 156, SEQ ID NO: 157, SEQ ID NO: 158, and SEQ ID NO: 159.
- 141. The method of claim 132, wherein said epidermal growth factor-like domain comprises an amino acid sequence set forth in SEQ ID NO: 152.
 - 142. The method of claim 132, wherein said polypeptide binds the p185erbB2 receptor.
 - 143. The method of claim 132, wherein said polypeptide is a recombinant polypeptide.



Creation date: 08-20-2004

Indexing Officer: TROBINSON - TERRI ROBINSON

Team: OIPEBackFileIndexing

Dossier: 08734592

Remarks:

Legal Date: 05-06-2002

| No. | Doccode | Number of pages |
|-----|---------|-----------------|
| 1 | A | 2 |
| 2 | CLM | 1 |
| 3 | REM | 3 |

| 3 | REM | 3 |
|--------------|--------------------|---|
| - | h 1 | |
| l otal r | number of pages: 6 | |

Order of re-scan issued on